


DETERMINING THE PRESENCE OF GLYPHOSATE-RESISTANT HORSEWEED UNDER FIELD CONDITIONS

Weed
Facts
WF-13



Glyphosate-resistant horseweed (or marestail or *Conyza canadensis*) has been confirmed throughout Delaware, Maryland and New Jersey. Horseweed is a winter annual weed. Emergence begins in the fall and continues into the early summer. The plant initially grows low to the ground (rosette stage) and then develops a tall erect stem (bolting stage), from which numerous seed heads are produced. Resistant-horseweed can tolerate glyphosate treatments at all stages, even while the plant is in the rosette stage.

Glyphosate is the active ingredient in Roundup, Touchdown, Duramax, and numerous other brand names.

If resistance is suspected in one of your fields, some questions should be asked to help determine if herbicide resistance is the reason for lack of control.

In trying to determine if herbicide resistant weeds are present, first eliminate other causes of herbicide failure.

- was herbicide applied as recommended?
- did the plants emerge after application?
- are there patterns to where weeds were not controlled in the field? (this implies a problem with application)



The lower (1) and upper (2) portions of a mature horseweed plant. Drawing from WEEDS OF THE NORTH CENTRAL STATES, University of Illinois Urbana-Champaign, Bulletin 773.

Resistance develops by selecting resistant-biotypes through repeated use of the same herbicide mode of action. Some questions regarding herbicide use need to be considered:

- has the same herbicide (or herbicides) with the same mode of action been used consistently over a period of years?
- has the species been controlled by the herbicide(s) in the recent past?
- has there been a decline in control with the suspected species?

One of the classic signs for resistance (for all situations where resistance is to a postemergence herbicide) is that only one weed species is present, all others are effectively controlled (or uniformly suppressed depending on the herbicide and weed).



The level of resistance in the horseweed populations is not as great as seen with other herbicide resistant weeds. It is common to see some glyphosate response in the resistant horseweed. The response can be either yellowing in the growing point or death of the growing point, resulting in long branches developing from lateral buds on the lower part of the plant. Given the situation in our region, there should be some susceptible plants (controlled by glyphosate) in the

same field along with resistant plants (surviving plants). This occurrence of controlled and uncontrolled plants randomly scattered throughout the field is an easy way to identify fields with resistant horseweed (see above photo).



Photo on the left is a plant sprayed with glyphosate and showing no difference in growth due to the treatment. Photo on the right is a horseweed plant treated with glyphosate and the top portion of the plant was killed, but the plant does not die and begins to produce long branches from lateral buds on the lower portion of the stem. Both plants are resistant to glyphosate.

If you suspect resistance, but are still uncertain or your situation is not covered by the situations described above, there is a test you can do yourself. Mix up a small hand sprayer with 1% solution of glyphosate (2.5 tablespoons per 1 gallon of water or one-half cup per 3 gallons of water) and spray the suspected plants. Be sure to check if a nonionic surfactant is required with the form of glyphosate you are using (if needed 2.5 tablespoons per 1 gallon of water or one-half cup per 3 gallons of water). Spray approximately 30 plants in a small area. Do not “douse” the plants with the spray mixture, rather a light spray is adequate. Then spray the rest of the field with the appropriate herbicide to prevent the horseweed from producing seed. This procedure will help you determine if glyphosate-resistant horseweed is present in your field, which in turn will allow better management decisions to be made. Once a field is identified with glyphosate-resistant horseweed, glyphosate will no longer control this species and additional management will be needed for the present year and many future years.

Mark VanGessel, Professor/Extension Specialist, Weed Science and Crop Management
 Quintin Johnson, Extension Associate III, Weed Science and Crop Management
 Department of Plant and Soil Sciences, University of Delaware

Cooperative Extension Education in Agriculture and Home Economics, University of Delaware, Delaware State University and the United States Department of Agriculture cooperating. Distributed in furtherance of Acts of Congress of May 8 and June 30, 1914. It is the policy of the Delaware Cooperative Extension System that no person shall be subjected to discrimination on the grounds of race, color, sex, disability, age, or national origin.

Commercial companies or products are mentioned in this publication solely for the purpose of providing specific information. Mention of a company or product does not constitute a guarantee or warranty of products by the Agricultural Experiment Station or Delaware Cooperative Extension or an endorsement over products of other companies not mentioned.